

Register of Australian Herbage Plant Cultivars

B. Legumes

11. Serradella

(a) *Ornithopus compressus* L. (yellow serradella) cv. Paros

Reg. No. B-11a-8

Registered 15 January 1990

Originators: J. S. Gladstones^A, M. D. A. Bolland^A,
C. K. Revell^B

^AWestern Australian Department of Agriculture, Baron-Hay Court, South Perth, W.A. 6151, Australia.

^BWestern Australian Department of Agriculture, Dryland Research Institute, P.O. Box 432, Merredin, W.A. 6415, Australia.

Registrar: R. N. Oram

CSIRO Division of Plant Industry, G.P.O. Box 1600, Canberra, A.C.T. 2601, Australia.

Released by Western Australian Department of Agriculture

Australian Journal of Experimental Agriculture, 1990, 30, 443–4.

Origin

Paros was collected by D. J. Gillespie in 1978, 2 km east of Parikia on the island of Paros, Greece. The collection site has an altitude of 100 m and an annual rainfall of approximately 475 mm. The site was a moderately grazed donkey track with a grey gritty sand soil type, stoney, pH 6.0. Paros was tested under the collector's code number DP6.

It was included in a 1981 distribution of 32 lines and a 1982 distribution of 100 selected serradella lines for testing across southern and eastern mainland Australia (Gladstones 1984).

Paros was submitted by the Western Australian Department of Agriculture and recommended for registration by the Western Australian Herbage Plant Liaison Committee. The Western Australian Department of Agriculture will maintain breeders' seed.

Morphological description

Paros differs from the 'typical' *Ornithopus compressus* cvv. Pitman and Tauro in the following characters: growth remains prostrate and spreading well into the flowering stage; stems and pods are more strongly pigmented, especially where exposed to the sun, pigment brownish purple; pods are non-segmented, much more strongly curved and do not break up readily into single-seeded segments at maturity. This contrasts to pods of Madeira which are segmented and break up under moderate mechanical stress, and pods of Pitman which are segmented and break up spontaneously after maturity.

The leaves of Paros have up to 12 leaflets pairs and are subtended by characteristically large and prominent bracts with 4.5–5.5 leaflet pairs; inflorescences with 2–5 flowers and pods; seeds up to 12 per pod. Seeds are a little larger than those of Pitman and Tauro, 250 000–300 000 seeds/kg dehulled.

Agronomic characters

Paros flowers early, about 93 days at Perth with mid-May sowing, that is, about 2 and 20 days earlier than cvv. Madeira and Tauro, respectively. However, the rate of pod maturation in Paros is more rapid than in Madeira, mature seed being produced approximately 6 days earlier (D. L. Lloyd, unpublished data).

Clean dehulled seed production of Paros has been impressive with highest yields being in the order of 800–900 kg/ha. In a group of 33 lines tested over 4 years at Esperance (W.A.), Paros consistently rated in the top 5 for seed production. At Albany (W.A.) in 1985, Paros was rated in the top 6 lines for seed production in two, Stage 2 evaluation trials. In both trials Paros was rated ahead of Madeira. In 1987, seed yield of Paros exceeded that of Madeira at both Merredin and Geraldton (W.A.), although total seed production at these sites was low.

Seed production of Paros in South Australia has also been consistently high, being second to Tauro at Stewarts Range and Conmurra in 1985, and second to Uniserra at Keith in 1985. At Stewarts Range in 1986, it was ranked 12 out of 26 lines but still produced 639 kg/ha of dehulled seed. In Queensland (D. L. Lloyd, unpublished data), seed production of Paros has been excellent, though slightly less than Madeira. In 1985 at Toowoomba, Paros produced 390 kg/ha clean, dehulled seed compared with 540 kg/ha from Madeira. In 1985 at Stanthorpe, Paros produced 2.72 t/ha podded seed compared with 2.86 t/ha from Madeira.

Paros is particularly hardseeded and laboratory tests show it to have a very slow rate of softening. This contrasts to Pitman which has a substantially lower hardseed content (Bolland 1987).

Regeneration of Paros in the second year has often been lower than Madeira, probably due to a slower rate of hardseed softening. This would also appear related to the non-segmented pod morphology of Paros, resulting in less pod–soil contact than occurs with the segmented Madeira. Dry matter production is often lower during early winter, although differences are not as great in spring. Regeneration after cereal cropping has, however, been particularly good, being superior to Pitman and Eneabba at Merredin (W.A.) and to Tauro at South Stirling Ranges, Dinninup and Arthur River (W.A.), in 1986.

Paros is tolerant of high aluminium concentrations, in contrast to Madeira which is particularly sensitive (Drew 1987).

Paros may be useful in mixtures with Madeira on sandy soils in low rainfall areas of Western Australia (300–375 mm annual average). Its high level of hardseed should improve long term persistence of serradella, particularly in situations

where the pasture is cropped. Its tolerance of high aluminium levels should give it an advantage over Madeira on strongly acid soils. It could also fill similar niches in central and northern New South Wales (R. D. Freebairn, pers. comm.). Paros will also be used in mixtures of serradella cultivars (particularly Madeira, Eneabba and Jebala) for sowing on sandy soils in hotter, drier areas of southern inland Queensland (D. L. Lloyd, pers. comm.).

Acknowledgments

Paros was tested as part of an interstate collaborative testing scheme for serradellas. The main field trials in Western Australia were conducted by M. D. A. Bolland (Esperance), L. Cransberg (Albany) and M. A. Ewing (Merredin); in South Australia by A. D. Craig (Kybybolite); in New South Wales by T. P. Drew (Trangie) and R. D. Freebairn (Coonabarabran); in Queensland by D. L. Lloyd (Toowoomba); and in Victoria by S. G. Clark (Hamilton) and L. J. Hamilton (DARA, Bairnsdale).

Pure seed production was supervised by D. A. Nicholas and C. J. Sykes (Western Australian Department of Agriculture). Funds were provided by the Herbage Seed Trust Fund.

The introduction and initial evaluation of Paros were assisted financially by the Wheat Industry Research Committee of Western Australia and the Australian Wool Research Trust Fund.

References

- Bolland M. D. A. (1987). Serradella, subterranean clover and medic research at Esperance, Western Australia. Western Australian Department of Agriculture, Technical Bulletin No. 76.
- Drew T. P. (1987). Serradella aluminium tolerance experiment. Report for New South Wales. Serradella Newsletter No. 1.
- Gladstones J. (1984). Selection of serradella genotypes for Australian agriculture. *Australian Plant Breeding and Genetics Newsletter* **34**, 58-62.